EURASIAN EXPERIMENT JOURNAL OF SCIENTIFIC AND APPLIED RESEARCH
(EEJSAR) ISSN: 2992-4146
©EEJSAR Publications Volume 7 Issue 3 2025

Phytoestrogens in Menopause: Mechanisms and Clinical Implications

Omeye Francis I.

Faculty of Medicine Kampala International University Uganda

ABSTRACT

Menopause is a natural physiological transition marked by the decline in estrogen levels, leading to various symptoms such as vasomotor instability, osteoporosis, cardiovascular risks, and cognitive decline. Phytoestrogens, plant-derived compounds with estrogen-like activity, have emerged as potential alternatives to hormone replacement therapy (HRT) for managing menopausal symptoms. This review explores the mechanisms of phytoestrogens, their influence on estrogen receptors, metabolism, and biological effects. The major classes of phytoestrogens, including isoflavones, lignans, and coumestans, are discussed, along with their sources and pharmacokinetics. Clinical studies evaluating the efficacy and safety of phytoestrogens in alleviating menopausal symptoms, improving bone density, and reducing cardiovascular risks are reviewed. Furthermore, potential risks and limitations, including bioavailability, individual variability in metabolism, and interactions with medications, are highlighted. While phytoestrogens show promise as a natural alternative for menopausal management, further large-scale clinical trials are necessary to establish optimal dosages and long-term safety.

Keywords: Phytoestrogens, Menopause, Isoflavones, Estrogen Receptors, Hormone Replacement Therapy

INTRODUCTION

Menopause is a natural biological transition that occurs in women, typically between the ages of 45 and 55, marking the cessation of ovarian function and the subsequent decline in circulating estrogen levels [1]. This hormonal shift triggers a cascade of physiological changes, affecting multiple organ systems and leading to symptoms such as hot flashes, night sweats, mood disturbances, osteoporosis, and an increased risk of cardiovascular diseases [2]. The decline in estrogen also has implications for metabolic health, cognition, and overall quality of life. Traditionally, hormone replacement therapy (HRT) has been the primary intervention for managing menopausal symptoms and preventing long-term complications associated with estrogen deficiency [3]. HRT has been shown to effectively alleviate vasomotor symptoms, preserve bone mineral density, and offer cardioprotective benefits. However, concerns regarding the potential risks of HRT, including an increased likelihood of breast cancer, thromboembolic events, and stroke, have led to a growing interest in alternative therapies [4,5]. Among these alternatives, phytoestrogens have gained considerable attention due to their structural similarity to endogenous estrogens and their ability to bind to estrogen receptors [6]. Phytoestrogens are naturally occurring plant-derived compounds classified into three main groups: isoflavones, lignans, and coumestans [7]. These compounds exhibit both estrogenic and anti-estrogenic properties, depending on endogenous estrogen levels and receptor subtype affinity. They are commonly found in foods such as soybeans, flaxseeds, whole grains, and legumes [8]. Several studies suggest that phytoestrogens may help alleviate menopausal symptoms, improve bone health, and support cardiovascular function, offering a potential alternative to conventional HRT [9]. However, variations in individual metabolism, differences in gut microbiota composition, and inconsistencies in clinical study outcomes highlight the need for further research to determine optimal dosages and long-term safety [10]. This review explores the underlying mechanisms of phytoestrogens, their clinical applications, and the potential benefits and risks associated with their use in menopausal management.

Mechanisms of Action of Phytoestrogens

Phytoestrogens are plant-derived polyphenolic compounds that exert estrogenic or anti-estrogenic effects by binding to estrogen receptors (ERs), particularly ER-α and ER-β. These compounds mimic or modulate estrogenic activity by competing with endogenous estrogens for receptor binding. Their effects depend on receptor affinity, tissue distribution, and endogenous hormone levels [11]. Isoflavones, primarily found in soybeans, function as

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

selective estrogen receptor modulators (SERMs) [12]. They preferentially bind to ER-β, which is more abundant in bone, cardiovascular, and brain tissues. This selective binding results in beneficial effects such as bone protection, improved lipid metabolism, and potential neuroprotection while minimizing risks associated with ER-α activation in breast and uterine tissues [13].Lignans, present in flaxseeds, undergo microbial metabolism in the gut, converting into enterolactone and enterodiol, which exhibit weak estrogenic activity. These metabolites interact with ERs and modulate estrogen metabolism, leading to potential protective effects against hormone-dependent cancers and cardiovascular diseases [14]. Coumestans, found in legumes like alfalfa and clover, exhibit strong estrogenic activity due to their high affinity for ERs. While their physiological significance in humans is less understood, they may contribute to modulating hormonal balance and reducing menopausal symptoms [15]. Beyond receptor binding, phytoestrogens influence gene expression, antioxidant activity, and anti-inflammatory pathways [16]. They modulate enzymatic activity involved in estrogen synthesis and metabolism, impacting overall hormonal balance. Their role in regulating oxidative stress and inflammation further supports their potential benefits in menopausal management. However, variations in bioavailability and individual metabolism impact their efficacy, necessitating further research to optimize clinical applications.

Clinical Implications of Phytoestrogens in Menopausal Management

Vasomotor symptoms, including hot flashes and night sweats, are among the most common menopausal complaints. Clinical studies suggest that isoflavones can alleviate these symptoms, although the extent of benefit varies based on individual metabolic differences and dietary intake. Some women experience significant symptom relief, while others show minimal response [17]. Bone health is a critical concern in menopause due to increased bone resorption and reduced bone mineral density. Phytoestrogens, particularly isoflavones and lignans, have demonstrated the ability to enhance bone density by influencing osteoblastic activity and inhibiting osteoclastic resorption [18]. Their role in osteoporosis prevention offers a potential alternative for women seeking non-hormonal interventions. Cardiovascular health is another significant area of concern, as estrogen deficiency increases the risk of atherosclerosis and hypertension. Phytoestrogens exert lipid-lowering effects, improve endothelial function, and exhibit vasodilatory properties. These factors contribute to reduced cardiovascular risks in menopausal women, making phytoestrogens a promising natural intervention [11]. Cognitive function and neuroprotection have also been linked to phytoestrogen consumption. Estrogen plays a key role in brain function, and its decline has been associated with cognitive decline and an increased risk of neurodegenerative diseases. Phytoestrogens exhibit antioxidant and anti-inflammatory properties that may support cognitive health, although more research is required to determine their long-term efficacy in preventing cognitive impairment [19].

Safety, Bioavailability, and Limitations

Phytoestrogens, plant-derived compounds with estrogen-like activity, offer potential health benefits, particularly in hormone-related conditions [12]. However, their effectiveness is influenced by several factors, including bioavailability, safety concerns, and possible drug interactions.

Bioavailability

The bioavailability of phytoestrogens varies significantly among individuals due to differences in gut microbiota composition [20]. The conversion of phytoestrogens into their biologically active forms depends on specific bacterial species in the intestines. Some individuals, known as equol producers, can metabolize soy isoflavones into equol, a more potent phytoestrogen with enhanced bioactivity [21]. Others lack the necessary gut bacteria, reducing the efficacy of phytoestrogen supplementation. Additionally, factors such as age, diet, and overall gut health play a role in absorption and metabolism, affecting the overall physiological response.

Safety Concerns

Although phytoestrogens are generally regarded as safe, their potential estrogenic effects raise concerns, particularly in individuals with hormone-sensitive conditions. Some studies suggest that phytoestrogens may stimulate the proliferation of estrogen-sensitive tissues, potentially influencing conditions such as breast and endometrial cancers [22]. However, research on their exact role remains inconclusive, with some evidence indicating a protective effect against hormone-related malignancies. Given these conflicting findings, caution is advised for individuals with a history of estrogen-sensitive cancers, and medical supervision is recommended before incorporating phytoestrogens into the diet [23].

Drug Interactions

Phytoestrogens can interact with various medications, potentially altering their efficacy. For instance, they may interfere with anticoagulant drugs, affecting blood clotting mechanisms [24]. Additionally, their estrogen-like properties may impact the effectiveness of hormone therapies, such as tamoxifen, used in breast cancer treatment. Individuals on these medications should consult healthcare professionals before using phytoestrogen supplements to prevent unintended interactions and ensure safe use [25,26]. While phytoestrogens offer promising health

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

benefits, understanding their bioavailability, safety considerations, and potential drug interactions is essential for informed use.

CONCLUSION

Phytoestrogens provide a natural alternative for managing menopausal symptoms, offering potential benefits in vasomotor regulation, bone health, cardiovascular protection, and cognitive function. Their estrogen-like activity may help alleviate hot flashes, support bone density, and improve lipid profiles. However, individual responses vary due to differences in metabolism and gut microbiota. Concerns about long-term safety, particularly in hormone-sensitive conditions, highlight the need for cautious use. Standardized dosages remain a challenge, as the optimal intake for efficacy and safety is not well established. Additionally, potential interactions with hormone therapies and anticoagulants require further investigation. Large-scale clinical trials are necessary to determine the long-term effects, establish appropriate dosages, and optimize formulations. Future research should focus on understanding the molecular mechanisms of phytoestrogens and their impact on menopausal health. Until then, medical supervision is recommended for individuals considering phytoestrogen supplementation as part of their menopausal management strategy.

REFERENCES

1. Abdi F, Rahnemaei FA, Roozbeh N, Pakzad R. Impact of phytoestrogens on treatment of urogenital menopause symptoms: A systematic review of randomized clinical trials. Eur J Obstet Gynecol Reprod Biol. 2021 Jun;261:222-235. doi: 10.1016/j.ejogrb.2021.03.039. Epub 2021 Apr 20. PMID: 33962824.
2. Franco OH, Chowdhury R, Troup J, Voortman T, Kunutsor S, Kavousi M, Oliver-Williams C, Muka T. Use of Plant-Based Therapies and Menopausal Symptoms: A Systematic Review and Meta-analysis. JAMA. 2016 Jun 21;315(23):2554-63. doi: 10.1001/jama.2016.8012. PMID: 27327802.
3. Dizavandi FR, Ghazanfarpour M, Roozbeh N, Kargarfard L, Khadivzadeh T, Dashti S. An overview of the phytoestrogen effect on vaginal health and dyspareunia in peri- and post-menopausal women. Post Reprod Health. 2019 Mar;25(1):11-20. doi: 10.1177/2053369118823365. Epub 2019 Feb 20. PMID: 30786797.
4. Chen MN, Lin CC, Liu CF. Efficacy of phytoestrogens for menopausal symptoms: a meta-analysis and systematic review. Climacteric. 2015 Apr;18(2):260-9. doi: 10.3109/13697137.2014.966241. Epub 2014 Dec 1. PMID: 25263312; PMCID: PMC4389700.
5. Lethaby A, Marjoribanks J, Kronenberg F, Roberts H, Eden J, Brown J. Phytoestrogens for menopausal vasomotor symptoms. Cochrane Database Syst Rev. 2013 Dec 10;2013(12):CD001395. doi: 10.1002/14651858.CD001395.pub4. PMID: 24323914; PMCID: PMC10247921.
6. Lipovac M, Chedraui P, Gruenhut C, Gocan A, Kurz C, Neuber B, et al. The effect of red clover isoflavone supplementation over vasomotor and menopausal symptoms in postmenopausal women. Gynecol Endocrinol. 2012 Mar;28(3):203-7.
7. Faure ED, Chantre P, Mares P. Effects of a standardized soy extract on hot flushes: a multicenter, double-blind, randomized, placebo-controlled study. Menopause. 2002 Sep-Oct;9(5):329-34.
8. Messina M, Mejia SB, Cassidy A, Duncan A, Kurzer M, Nagel SC, et al. Neither soyfoods nor isoflavones warrant classification as endocrine disruptors: A technical review of the observational and clinical data. Crit Rev Food Sci Nutr. 2021 Mar 27;61(4):537-553.
9. Nassan FL, Chavarro JE, Tanrikut C. Diet and men's fertility: does diet affect sperm quality? Fertil Steril. 2018 Sep;110(4):570-577.
10. Messina M, Rogero MM, Fisberg M, Waitzberg D. Health impact of childhood and adolescent soy consumption. Nutr Rev. 2017 Jan;75(1):500-515.
11. Mense SM, Hei TK, Ganju RK, Bhat HK. Phytoestrogens and breast cancer prevention: possible mechanisms of action. Environ Health Perspect. 2008 Apr;116(4):426-33. doi: 10.1289/ehp.10538. PMID: 18414622; PMCID: PMC2291001.
12. Messina MJ, Wood CE. Soy isoflavones, estrogen therapy, and breast cancer risk: analysis and commentary. Nutr J. 2008 Jun 3;7:17. doi: 10.1186/1475-2891-7-17. PMID: 18522734; PMCID: PMC2443803.
13. Song S, Moon HG, Noh DY, Jung SY, Lee ES, Kim Z, Youn HJ, Cho J, Yoo YB, Lee SK, Lee JE, Nam SJ, Kim YS, Min JW, Jun S, Joung H, Lee JE. Dietary isoflavone intake among breast cancer survivors and cancer-free women. Nutr Res Pract. 2025 Feb;19(1):80-95. https://doi.org/10.4162/nrp.2025.19.1.80
14. Corona, G., Kreimes, A., Barone, M. *et al.* Impact of lignans in oilseed mix on gut microbiome composition and enterolignan production in younger healthy and premenopausal women: an in vitro pilot study. *Microb Cell Fact* **19**, 82 (2020). https://doi.org/10.1186/s12934-020-01341-0
15. Jeff L. Staudinger, Clinical applications of small molecule inhibitors of Pregnane X receptor, Molecular and Cellular Endocrinology,2019, 485, 61-71. https://doi.org/10.1016/j.mce.2019.02.002.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

16. Desmawati D, Sulastri D. Phytoestrogens and Their Health Effect. Open Access Maced J Med Sci. 2019 Feb 14;7(3):495-499. doi: 10.3889/oamjms.2019.044. PMID: 30834024; PMCID: PMC6390141.
17. Avis NE, Crawford SL, Green R. Vasomotor Symptoms Across the Menopause Transition: Differences Among Women. Obstet Gynecol Clin North Am. 2018 Dec;45(4):629-640. doi: 10.1016/j.ogc.2018.07.005. Epub 2018 Oct 25. PMID: 30401547; PMCID: PMC6226273.
18. Anastasia Armeni, Panagiotis Anagnostis, Eleni Armeni, Nikoletta Mili, Dimitrios Goulis, Irene Lambrinoudaki, Vasomotor symptoms and risk of cardiovascular disease in peri- and postmenopausal women: A systematic review and meta-analysis, Maturitas, 2023; 171, 13-20. https://doi.org/10.1016/j.maturitas.2023.02.004.
19. Soni M, Rahardjo TB, Soekardi R, Sulistyowati Y, Lestariningsih, Yesufu-Udechuku A, Irsan A, Hogervorst E. Phytoestrogens and cognitive function: a review. Maturitas. 2014 Mar;77(3):209-20. doi: 10.1016/j.maturitas.2013.12.010. Epub 2014 Jan 8. PMID: 24486046.
20. Viggiani MT, Polimeno L, Di Leo A, Barone M. Phytoestrogens: Dietary Intake, Bioavailability, and Protective Mechanisms against Colorectal Neoproliferative Lesions. Nutrients. 2019 Jul 24;11(8):1709. doi: 10.3390/nu11081709. PMID: 31344966; PMCID: PMC6722977.
21. Esma Karahmet Farhat, Emina Karahmet Sher, Amina Džidić-Krivić, Ines Banjari, Farooq Sher, Functional biotransformation of phytoestrogens by gut microbiota with impact on cancer treatment, The Journal of Nutritional Biochemistry, 2023; 118,109368. https://doi.org/10.1016/j.jnutbio.2023.109368.
22. Lv, J., Jin, S., Zhang, Y. *et al.* Equol: a metabolite of gut microbiota with potential antitumor effects. *Gut Pathog* **16**, 35 (2024). https://doi.org/10.1186/s13099-024-00625-9
23. Ahkam Saddam Seyed Hameed, Parkash Singh Rawat, Xiangfeng Meng, Weifeng Liu, Biotransformation of dietary phytoestrogens by gut microbes: A review on bidirectional interaction between phytoestrogen metabolism and gut microbiota, Biotechnology Advances,2020; 43,107576. https://doi.org/10.1016/j.biotechadv.2020.107576.
24. Mirmohammadali SN, Rosenkranz SK. Dietary phytochemicals, gut microbiota composition, and health outcomes in human and animal models. Biosci Microbiota Food Health. 2023;42(3):152-171. doi: 10.12938/bmfh.2022-078. Epub 2023 Apr 25. PMID: 37404568; PMCID: PMC10315191.
25. Gaya P, Medina M, Sánchez-Jiménez A, Landete JM. Phytoestrogen Metabolism by Adult Human Gut Microbiota. *Molecules*. 2016; 21(8):1034. https://doi.org/10.3390/molecules21081034
26. Williams CLYbarra AR, Meredith AN, Durrant BS, Tubbs CW. Gut Microbiota and Phytoestrogen-Associated Infertility in Southern White Rhinoceros. 2019; mBio10:10.1128/mbio.00311-19.https://doi.org/10.1128/mbio.00311-19

CITE AS: Omeye Francis I. (2025). Phytoestrogens in Menopause: Mechanisms and Clinical Implications. EURASIAN EXPERIMENT JOURNAL OF SCIENTIFIC AND APPLIED RESEARCH, 7(3):22-25

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited